

PERpose Plus™

Broad Spectrum Algacide and Fungicide

For use in Horticulture, Agriculture, and Natural Turf Industries

PERpose Plus™ is a treatment for the prevention and suppression/control of horticultural and agricultural diseases in commercial greenhouses, garden centers, landscapes, agricultural crops, nurseries, interiorscapes and professional turf care.

ACTIVE INGREDIENT:

Hydrogen Peroxide/Hydrogen Dioxide..... 33.00%
INERT INGREDIENTS: 67.00%
Total.....100.00%



Keep Out of Reach of Children

DANGER-PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

If in eyes:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15 to 20 minutes.Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment adviceHave person sip a glass of water if able to swallow.Do not induce vomiting unless told to do so by a poison control center or doctor.Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. To contact a regional poison center call 1-800-222-1222.

IN EMERGENCIAS CALL CHEMTREC: (800) 424-9300

PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS AND DOMESTIC ANIMALS

Danger. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed. **PERpose Plus™** may be fatal if absorbed through the skin. Do not get in eyes, on skin or clothing. Do not breathe vapor from concentrate. Wear protective eyewear (goggles, face shield, or safety glasses). Wear coveralls over long sleeved shirt and long pants, socks, chemical-resistant footwear and chemical-resistant gloves. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, mammals, fish and aquatic life. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA. This product is highly toxic to bees and other beneficial insects exposed to direct contact on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Do not apply this product or allow it to drift to crops where beneficial insects are part of an Integrated Pest Management strategy.

PHYSICAL AND CHEMICAL HAZARDS

PERpose Plus™ is a strong oxidizing agent. Mix only with water. PERpose Plus™ is not combustible; however, at temperatures exceeding 156°F, decomposition releases oxygen. The oxygen released could initiate or promote combustion of other materials.

Procedure for Leak or Spill - Stop leak if this can be done without risk. Shut off ignition sources; no flames, smoking, flares, or spark producing tools. Keep combustible and organic materials away. Flush spilled material with large quantities of water. Undiluted material should not enter confined spaces.

Personal Protective Equipment (PPE)

Applicators and handlers must wear coveralls over long sleeved shirt, long pants and chemical resistant foot wear, plus socks. When mixing and loading, wear a chemical resistant apron. For overhead exposure, wear chemical resistant headgear. Also wear protective eyewear (goggles, face shield, or safety glasses), and chemical resistant gloves. When cleaning equipment, wear a chemical resistant apron. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water.

User Safety Recommendations

User should wash hands thoroughly with soap and water before eating, drinking, using tobacco, or using the toilet. Users should remove clothing immediately if contaminated by pesticide. Wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Manufactured for: A Growing Alternative, Inc.
751 Fairview Church Road
Spruce Pine, NC 28777
828-766-6179

EPA Registration No. 86729-1

EPA Est. No. 5568-SC-001

Net Contents: 5 Gal-18.93L 15 Gal-56.78L 30 Gal-113.56 55 gal.-202.38L 275 Gal -1011.90 L

PERpose Plus™

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with labeling. Do not apply this product in a way that will contact workers, or other persons, either directly, or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for the pesticide regulation.

AGRICULTURE USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE), notification to workers and Restricted-Entry interval (REI). The requirements in this box only apply to the uses of this product that are covered by the Worker Protection Standard.

Except as indicated below, there is a restricted entry interval (REI) of zero (0) hours for Perpose Plus™. For interior fogging and spraying applications, there is a restricted entry interval of one (1) hour. For field applications, do not allow reentry until sprays have dried. Early reentry requires coveralls over long-sleeved shirt and pants, shoes with socks and waterproof gloves.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (30 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep unprotected persons out of treated areas until sprays have dried.

PERpose Plus™ is a treatment for the prevention and suppression/control of Horticultural, Agricultural and Turf diseases, in the following locations:

Commercial greenhouses	Garden centers	Nurseries	Professional turf care	Interiorscapes
Orchards	Field crops	Vineyards	Groves	Landscapes

PERpose Plus™ Treats, Controls and Prevents the following diseases:

<i>Algae</i>	<i>Aphanomyces</i>	<i>Anthracnose</i>	<i>Alternaria</i>
<i>Downy Mildew</i>	<i>Botrytis (grey mold)</i>	<i>Black Spot</i>	<i>Fusarium (root rot)</i>
<i>Erwinia</i>	<i>Phytophthora (blights, rots)</i>	<i>Leaf Spot</i>	<i>Powdery Mildew</i>
<i>Plasmopara</i>	<i>Pythium</i>	<i>Pseudomonas</i>	<i>Rust</i>
<i>Smut</i>	<i>Scab</i>	<i>Rhizoctonia</i>	<i>Xanthomonas</i>
<i>Wilts & Blights</i>	<i>Thielaviopsis</i>	<i>Uncinula (powdery mildew)</i>	<i>Sclerotinia</i>

PERpose Plus™ may be used on:

Ornamentals and Agriculture crops, including vegetable crops, fruit crops, tobacco, etc	Nursery stock Trees	Seeds and seedbeds Poinsettias	Seedlings Roses
Turf	Bedding plants	Cut flowers	Flowering plants
			Bulbs
			Cuttings

Compatibility

It is advisable to test a small sample of **PERpose Plus™** on plants prior to treating large numbers, as not all plant species have been tested.

Reference regarding usage rates for both Horticulture and Agriculture Applications

The rates listed for application of **PERpose Plus™**, are universal for both agriculture and horticulture use. The rates using **PERpose Plus™** for curative results are 1:100, or 1fl. oz. per gallon of clean water and 1/3 -1/4 fl. oz. per gallon of clean water used as a weekly preventative rate.

Using PERpose Plus™ for curative measures

When using **PERpose Plus™** curative rates, apply for 3 consecutive days drenching the entire plant, including leaves, stems, stalks and the soil surface to saturation. Afterward, maintain with the preventative rate on a weekly basis for best results.

PERpose Plus™ controls on contact

It is not advisable to apply PERpose Plus™ in dry, extremely hot conditions.

PERpose Plus™ concentrate must not be combined or mixed with any other pesticide or fertilizer. A diluted solution of **PERpose Plus™** may be used in combination with most soluble fertilizers, although a compatibility test should be completed prior to mixing a final batch.

Solution Preparation

PERpose Plus™ works best when diluted with water containing low levels of organic or inorganic materials and having a neutral PH. Rinse mixing tank thoroughly with water before mixing concentrate, residues may cause product to lose effectiveness.

PERpose Plus™ will readily mix with clean, neutral water and does not require additional agitation. Prepare a solution by filling the tank with the required volume of water and then add concentrate as required.

Do not reuse already mixed solution; make fresh daily.

The By-Product of **PERpose Plus™**, after application, is oxygen and water.

PERpose Plus™ is not compatible and may react with residues of recent foliar applications of metal based fungicides or supplements. For best results, apply **PERpose Plus™**, allow to dry, prior to application of metal based fungicides or supplements.

Non-metal based fungicides may be used in combination with PERpose Plus™. Please contact us for compatibility list.

PERpose Plus™ is effective via direct surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wetted. When used in accordance with label directions, diluted **PERpose Plus™** does not produce any visible residue or deleterious effects to plant. Leaf burn may occur if **PERpose Plus™** is used at higher than recommended application rates.

Do not apply this product through any irrigation system unless directed by the label. Refer to Chemigation Use Directions section.

For use on Surfaces and Equipment

PERpose Plus™ may be used as a fungicide and algicide in conjunction with a foaming adjuvant on the following sites:

Greenhouse structures	Benches	Watering systems
Pots	Evaporative coolers	Storage rooms
Ventilation equipment	Floors	Other equipment, etc

Conversion Table

fluid oz	milliliters	fluid oz	milliliters
1/4	7.4	2 1/2	74
1/3	9.9	4 1/3	128
1/2	15	5 1/3	158
3/4	22	6 1/4	185
1	30	10	296
1 2/3	49	23	680
2	59	128 (1gal.)	3785

Use **PERpose Plus™** to suppress/control fungi and slime forming algae on greenhouse structures including: glazing, plastic, benches, walkways, floors, walls, fan blades, coolers, storage rooms, ventilation ducts, watering systems, vats, tanks, bins, elevators, storage areas, food contact surfaces, process water systems, trucks and related equipment, conveyors, and spray equipment. Any equipment or structures pertaining to food contact surfaces should be rinsed with potable water after treatment with **PERpose Plus™**.

1. Sweep and remove all gross plant debris. Use power sprayer to remove loose dirt.
2. Use a dilution of 1:50 or 2.0 fl. oz. **PERpose Plus™** per gallon of clean water for surfaces that have not been pre-cleaned with water to remove organic deposits. A surfactant may be added if needed.
3. Apply solution in a manner that all surfaces are thoroughly wet. Acceptable application methods are: mop, sponge, power sprayer, or misting.
4. Heavy growths of algae and fungi may have to be manually removed following application. Use a solution of **PERpose Plus™** to wash away dead growth.
5. Reapply as often as needed to control algae and fungi.

Foaming application

Apply PERpose Plus™ in combination with a foaming agent on porous, irregular and steel surfaces until completely covered. Allow to dry and do not rinse. Add foaming agent only into dilute **PERpose Plus™** solution.

For evaporative coolers

Treat existing algae and slime contaminated surfaces with 1 fl. oz. **PERpose Plus™** per gallon of cooler water. Treat cooler water every week with a dilution rate of 1/3 fl. oz. to 1/4 fl. oz. **PERpose Plus™** per gallon of cooler water.

For irrigation systems including flooded floors, flooded benches, recycled water systems, humidification and misting systems.

1. Treat contaminated water with a dilution of 1/3 fl. oz. to 1/4 fl. oz. **PERpose Plus™** per gallon of water.

2. Treat clean water with a dilution of 1:10,000, or 1 gallon of **PERpose Plus™** concentrate per 10,000 gallons of water.

Foot Bath Mats

Using a dilution of 3/4 fl. oz. **PERpose Plus™** per gallon of water, fill foot bath mat to capacity. Use fresh solution daily.

Cutting Tools

Use a dilution of 1:100 - 1:300 or 1.0 fl. oz. to 1/3 fl. oz. **PERpose Plus™** per gallon of clean water. Drench tools to ensure full coverage. Additional surfactant may be used.

Flats, Trays and Pots

Use a dilution of 1:100 to 1:300 or 1.0 fl. oz. to 1/3 fl. oz. **PERpose Plus™** per gallon of clean water. Spray until runoff. Additional surfactant may be used.

Reference regarding usage rates for both Horticulture and Agriculture Applications

The rates listed for application of **PERpose Plus™**, are universal for both agriculture and horticulture use. The rates using **PERpose Plus™** for curative results are 1:100 or 1 fl. oz. per gallon of clean water and 1/3 -1/4 fl. oz. per gallon of clean water used as a weekly preventative rate.

Using PERpose Plus™ for curative measures

When using **PERpose Plus™** curative rates, apply for 3 consecutive days drenching the entire plant, including leaves, stems, stalks and the soil surface to saturation. Afterward, maintain with the preventative rate on a weekly basis for best results.

Mist Propagation of Cuttings and Plugs for both Horticulture and Agriculture Applications

Inject **PERpose Plus™** into the misting system to control/suppress algal and fungal disease, before disease becomes established on plant material. Inject **PERpose Plus™** using a dilution rate of 1:1000 on a consecutive basis for four to ten days. Reduce concentration to 1:5000 and continue application throughout the propagation cycle. At the first sign of disease, adjust concentration to 1:1000 or 1 gallon of **PERpose Plus™** per 1000 gallons of water.

Pre-plant dip treatment for both Horticulture and Agriculture Applications

Use **PERpose Plus™** in agriculture and ornamental applications, including nursery plants, seed beds, seeds, seedlings, bulbs or cuttings for the control/suppression of damping-off, root and other stem rot diseases caused by *Pythium*, *Phytophthora*, *Rhizoctonia*, *Botrytis*, *Crown Rot Powdery Mildew*, *Fusarium* or *Thielaviopsis*.

1. Use a dilution rate of 1/3 fl. oz. **PERpose Plus™** per gallon of water.
2. Immerse plants, or cuttings, remove and allow to drain. Do not rinse.

Soil Drench for Horticulture, Agriculture and Turf Application

PERpose Plus™ is effective for the control/suppression of *Pythium*, *Phytophthora*, *Rhizoctonia*, *Fusarium*, *Thielaviopsis* and other plant diseases. Use as a soil drench at the time of seeding or transplanting, as well as a periodic drench. **PERpose Plus™** can also be used on potting soil and growing mediums prior to planting.

1. Use a dilution of 1 fl. oz. per gallon of clean water.
2. Apply to saturation on soil or other growing media.

Foliar Spray Treatment in Greenhouses for both Horticulture and Agriculture Applications

PERpose Plus™ immediately controls/suppresses plant pathogens upon surface contact. Apply **PERpose Plus™** to horticulture ornamentals, agricultural crops, bedding plants, flowering plants, shrubs, trees and Turf. Apply solution to ensure thorough coverage and wetting of the foliage.

Initial (Curative) Application

1. Use a dilution of 1 fl. oz. **PERpose Plus™** per gallon of clean water. Make fresh daily; do not re-use already mixed solution.
2. Spray, or mist plants in early morning, or late evening. Ensure that all plant surfaces, including upper and lower foliage, stems, branches and stalks are fully wet with use solution.
3. Apply for one to three consecutive days. Follow directions for preventive treatment after the initial application.

Weekly (Preventative) Treatment:

1. Use a dilution of 1/3 fl. oz. **PERpose Plus™** per gallon of clean water.
2. Apply solution to plants by spray, mist, or fog.
3. Ensure that all plant surfaces, including upper and lower foliage, stems, branches and stalks are fully wet with use solutions. Spray every five to seven days as a preventative treatment. At the first signs of disease, spray daily with 1 fl. oz. of **PERpose Plus™** per gallon of water. Treat plants for three consecutive days and then resume weekly preventative treatment.

Foliar spray application in the field for Horticulture and Agriculture

PERpose Plus™ controls/suppresses disease immediately upon contact with any plant surface. Ensure that all plant surfaces, including upper and lower foliage, stems, branches and stalks are fully wet with use solution.

Initial (Curative) Application:

1. Use a dilution of 1 fl. oz. **PERpose Plus™** per gallon of clean water. Make fresh daily; do not re-use already mixed solution.
2. Spray or mist plants in early morning or late evening. Ensure that all plant surfaces, including upper and lower foliage, stems, branches and stalks are fully wet with use solution.
3. Apply for one to three consecutive days. Follow directions for preventive treatment after the initial application.

Weekly (Preventative) Treatment:

1. Use a dilution of 1/3 fl. oz. **PERpose Plus™** per gallon of clean water.
2. Apply solution to plants by spray, mist, or fog.
3. Ensure that all plant surfaces, including upper and lower foliage, stems, branches and stalks are fully wet with use solutions. Spray every five to seven days as a preventative treatment. At the first signs of disease, spray daily with 1 fl. oz. of **PERpose Plus™** per gallon of water. Treat plants for three consecutive days and then resume weekly preventative treatment.

Nursery stock and cut flowers for both Horticulture and Agriculture Applications

Apply **PERpose Plus™** to nursery stock such as:

Woody ornamentals	Bedding plants	Flowering plants
Roses	Container plants	Rhododendrons
Conifers	Azaleas	Shade trees
Agricultural crops		Natural Turf

Cut flowers

Use **PERpose Plus™** to prevent *Botrytis*, *Downy Mildew* and *Powdery Mildew* on flowers in transit or cold storage. Apply post-harvest.

Use 1/4 fl. oz. of **PERpose Plus™** per gallon of clean water. Spray flowers after grading and prior to storage or shipment. Repeat weekly for flowers in storage.

Bare root nursery stock for both Horticulture and Agriculture Applications

Use **PERpose Plus™** to prevent *Botrytis* on bud wood and nursery stock in storage.

Use a dilution of 1/4 -1/3 fl. oz. **PERpose Plus™** per gallon of water. Apply use dilution until plants are dripping wet or apply by dip method. Repeat weekly if necessary.

For seedbed treatment for Horticulture , Agriculture and Turf Applications

1. Prior to sowing seed, use a dilution of 2 fl. oz. **PERpose Plus™** per gallon of clean water.
2. Thoroughly wet or drench the seed bed to the point of saturation, with 60 to 100 gallons of dilute solution per 1000 square feet.
3. Let sit for one hour then immediately seed soil.
4. After seeds have germinated, use a dilution of 1 fl. oz. **PERpose Plus™** per gallon of clean water.
5. Wet soil and seedlings to a point of thorough wetness using light spray, or irrigation techniques.
6. Repeat application once per week until seed is well established.

For Use as a soil treatment prior to inoculation with beneficial microorganisms for Horticulture, Agriculture and Turf Applications

Use **PERpose Plus™** to reduce the number of pathogenic plant microorganisms in the soil.

1. Use a dilution of 2 fl. oz. **PERpose Plus™** per gallon of clean water.
2. Thoroughly wet, or drench the area to be inoculated.
3. Wait one day before inoculating the soil.

For turf treatment following inoculation of soil with beneficial microorganisms for Horticulture, Agriculture and Turf

Use **PERpose Plus™** to control plant pathogens on the foliar portion of turf.

Use a dilution 1 fl. oz **PERpose Plus™** per gallon of clean water. Apply to the turf by gently spraying leaf surfaces. Do not allow solution to be drenched into the soil and root systems, as a temporary reduction of beneficial soil microorganisms may occur.

Turf Applications

Broad spectrum treatment for control of algae, fungi on natural turf

PERpose Plus™ can be injected through automatic irrigation systems in turf areas. Refer to "Chemigation Directions" for specific instructions on using this product through irrigation systems.

Use **PERpose Plus™** to control *Anthraxnose*, *Brown Spot*, *Dollar Spot*, *Copper Spot*, *Fairy Ring*, *Pink Snow Mold*, *Pythium*, *Phytophthora*, *Summer Patch*, *Rhizoctonia*, *Scum*, *Fusarium Blight*, *Stripe Smut*, *Leaf Spot*, *Algae*, *Slime Molds* and their spores.

For use on all turf types including :

Commercial turf	Lawns	Athletic fields
Golf course fairways	Greens and tees	

On the following grasses:

Bermuda grass	Fescue	Rye Grass
Blue grass	Bent grass	St. Augustine including turf combinations

For Preventative Treatment of Algae/Fungus on Natural Turf

Typical treatment rates involve treating approximately 1000 square feet of lawn area with 1 to 10 gallons of diluted solution. Spray entire area until saturation to ensure the solution penetrates algal crusts and deposits. For best results use a spreader surfactant.

Initial (curative) Treatment

For treatment of heavy infestations of algae, dilute 5 1/3 to 10 fl. oz. of **PERpose Plus™** in 5 gallons of clean water. Apply 5-10 gallons of dilute solution per 1000 square feet.

Preventative Treatment

To prevent algal diseases, dilute 1 2/3, to 4 1/3 fl. oz. of **PERpose Plus™** in 5-10 gallons of clean water. Use 1-5 gallons of diluted solution per 1000 square feet.

1. Applications may be repeated every 5 to 7 days, or as necessary to control disease.
2. For best results, apply in early morning, or late afternoon.
3. Apply immediately after grass has been cut.
4. Applications may also be made during wet or rainy weather.

For severe conditions of crusted algae

1. Dilute 23 fluid ounces of **PERpose Plus™** per 2-10 gallons of clean water.
2. Apply 2-10 gallons of use solution to 1000 square feet of affected area.
3. Increased rates of active ingredient and increases in water volume may be required to penetrate layers of algae under severe conditions.
4. Under severe conditions, applications may be doubled by either increasing the amount of active ingredient per 1000 square feet of turf, or by applying treatment solution twice over the same area.

For fungicide treatment of natural turf

For initial (curative) treatment of heavy infestations of fungal disease

1. Dilute 5 1/3 to 10 fl. oz. **PERpose Plus™** in 5 gallons of clean water. For every 1000 square feet of turf, use 5-10 gallons of dilute solution.
2. Spray entire area until run-off. A spreader surfactant may be used for enhanced results.
3. For best results, apply uniformly over the area immediately after grass has been cut.

For preventative treatment of fungal disease

1. Dilute 2 1/2 to 6 1/4 fl. oz. **PERpose Plus™** per 5-10 gallons of clean water. Apply 5-10 gallons of dilute solution per 1000 square feet.
2. Spray entire area until run-off. A spreader surfactant may be used for enhanced results.
3. Start applications at the first sign of disease and repeat every 5 to 7 days, or as needed to control new, or established disease conditions.
4. For best results, apply uniformly over the area immediately after grass has been cut.
5. Applications may be repeated every 5 to 7 days, or as necessary to control disease.
6. For best results, apply in early morning, or late afternoon.
7. Apply immediately after grass has been cut.
8. Applications may also be made during wet or rainy weather.

Chemigation Directions for Use

General Requirements:

1. Apply this product through one of the following types of irrigation systems: center pivot, lateral move, end tow, side wheel roll, traveler, solid set, hand move, flood basin, or drip trickle irrigation system. Do not apply this product through any other type of irrigation system.
2. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
3. Ensure that the irrigation system is properly calibrated. If you have questions about calibration, contact the State Extension Service specialist, the equipment manufacturer, or other experts.
4. Do not connect irrigation systems (including greenhouse systems) used for pesticide application to a public water system unless proper safety devices for public water systems are in place. Read specific requirements provided below.
5. A person knowledgeable of the chemigation system and responsible for operation, or under the supervision of the responsible person, shall shut the system down and make any necessary adjustments should the need arise.

Specific Requirements for all Irrigation Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days throughout the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top of overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of liquid back towards the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump or equivalent, effectively designed and constructed of materials that are compatible with pesticides and capable of being filled with a system interlock.
7. Do not apply outdoors when wind speed favors drift beyond the area intended for treatment.

Application Instructions:

1. Remove scale, pesticide residues and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residue may cause product to lose effectiveness of strength.
2. Determine the treatment rates as indicated in the directions for use and make proper dilutions.
3. Prepare a solution by filling the tank with the required volume of water and then adding product as required. The product will immediately go into solution without any required agitation.
4. **PERpose Plus™** must not be applied in conjunction with any other pesticides or fertilizers as this may reduce performance of the product and should be avoided.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in the original container in a cool area out of direct sunlight and away from combustibles.

PESTICIDE DISPOSAL: Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance .

CONTAINER DISPOSAL; Non-refillable container up to 5 gallons. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Empty rinsate into application equipment, a mix tank, or store rinsate for later use, or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available. Non-refillable container greater than 5 gallons. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Turn the container onto its other end and tip it back and forth several times. Empty rinsate into application equipment, a mix tank, or store rinsate for later use, or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

Warranty

This material conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing, method of application, weather, watering practices, nature of soil, potting medium, disease problem, the condition of crop, incompatibility with other chemicals, pre-existing conditions and other conditions influencing the use of this product are beyond the control of the seller. Buyer assumes all risks associated with the use, storage, or handling of this material not in strict accordance with directions given herewith. TO THE EXTENT ALLOWABLE BY STATE LAW, NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.